

Multiplicities of representations in spaces of modular forms

BinYong Hsie

LMAM, Department of Mathematics, PeKing University,
BeiJing, P.R.China 100871
e-mail: byhsie@math.pku.edu.cn

Abstract

This paper shows that for a given irreducible representation ρ of Γ/Γ_1 , the two functions $\dim(M_k(\Gamma_1, \rho))$ and $\dim(S_k(\Gamma_1, \rho))$ of k are almost linear functions.

Let Γ and Γ_1 be two Fuchsian subgroups of $SL_2(\mathbb{Q})$ of the first kind, with Γ_1 a normal subgroup of Γ of index μ . Let $\mathbb{H}^*$ denote the extended upper half-plane [4] and put $Y = \Gamma \backslash \mathbb{H}^*$ and $X = \Gamma_1 \backslash \mathbb{H}^*$. Let $A_k(*)$ denotes the space of meromorphic modular forms for $*$ of weight k . We define a representation π_k of Γ on $A_k(\Gamma_1)$ by the formula

$$\pi_k(\gamma\Gamma_1)(f) = f|[\gamma^{-1}]_k \quad (\gamma \in \Gamma, \quad f \in A_k(\Gamma_1)),$$

where the notation $[*]_k$ is as in Shimura [4]. The representation π_k factors through Γ/Γ_1 to give a representation – also denoted π_k – of the latter group. The space $M_k(\Gamma_1)$ of holomorphic modular forms of weight k for Γ_1 and the subspace $S_k(\Gamma_1)$ of cusp forms of weight k for Γ_1 are both stable under π_k , and the resulting representations of Γ/Γ_1 on $M_k(\Gamma_1)$ and $S_k(\Gamma_1)$ will be denoted ρ_k and σ_k respectively.

Henceforth ρ denotes an irreducible complex representation of Γ/Γ_1 . If $-I \in \Gamma$ then the value of ρ on the coset of $-I$ in Γ/Γ_1 is a scalar by Schur's lemma, and we say that ρ is *even* or *odd* according as the scalar is 1 or -1 . If $-I \in \Gamma_1$ then ρ is automatically even.

If π is any finite-dimensional complex representation of Γ/Γ_1 then we write $\langle \rho, \pi \rangle$ for the multiplicity of ρ in π . For example $\langle \rho, \rho_{reg} \rangle = \dim \rho$, where ρ_{reg} is the regular representation of Γ/Γ_1 .

Theorem *Fix an irreducible complex representation ρ of Γ/Γ_1 , and put*

$$c = \frac{1}{4\pi} \int_{\Gamma \backslash \mathbb{H}} \frac{dx dy}{y^2}.$$

If $-I \notin \Gamma$ then

$$\lim_{k \rightarrow \infty} \frac{\langle \rho, \rho_k \rangle}{k \langle \rho, \rho_{reg} \rangle} = \lim_{k \rightarrow \infty} \frac{\langle \rho, \sigma_k \rangle}{k \langle \rho, \rho_{reg} \rangle} = c.$$

⁰Mathematical Subject Classification: 11F11. Key Words: modular form, cusp form.

If $-I \in \Gamma$ then the same assertion holds provided k runs through positive integers of the same parity as ρ .

Proof. We shall prove our assertion only for ρ_k . The proof for σ_k is similar.

For any positive integer k , let $i(k) = (-1)^k$. If $-I \in \Gamma_1$, we assume k is always even.

Let $\mathcal{S}$ be the set of the irreducible representations of Γ/Γ_1 with the same parity as k if $-I \in \Gamma$, and the set of all irreducible representations of Γ/Γ_1 otherwise.

Let p be a non-cusp point of Y which has μ different liftings $p_1, \dots, p_\mu$ on X , and let $\tilde{p}_j$ be a lifting of p_j in $\mathbb{H}^*$ ($1 \leq j \leq \mu$). By Riemann-Roch theorem we know that there exists an $f \in A_{i(k)}(\Gamma_1)$ such that f has poles only at cusps and $f(\tilde{p}_1) = 1$, $f(\tilde{p}_j) = 0$ for all $j = 2, \dots, \mu$. For any $\alpha \in \Gamma/\Gamma_1$, put $f_\alpha = f|[\alpha]_{i(k)}$. For a fixed sufficiently large even integer n_0 , we may choose $\varphi = \varphi_{n_0} \in S_{n_0}(\Gamma)$ such that φf_α ($\alpha \in \Gamma/\Gamma_1$) are all cusp forms. Let W denote the $\mathbb{C}$ -linear subspace of $M_{i(k)+n_0}(\Gamma_1)$ spanned by $\{\varphi f_\alpha | \alpha \in \Gamma/\Gamma_1\}$. Clearly, W is stable under $\pi_{i(k)+n_0}$. It is easy to see that

$$W \cong \bigoplus_{\rho \in \mathcal{S}} \langle \rho, \rho_{\text{reg}} \rangle \rho$$

since the two hand sides have the same trace.

When $k \geq i(k) + n_0$, the choice of f ensures that the natural map

$$W \bigotimes_{\mathbb{C}} M_{k-i(k)-n_0}(\Gamma) \rightarrow M_k(\Gamma_1)$$

is injective. Hence

$$\frac{\langle \rho, \rho_k \rangle}{k \langle \rho, \rho_{\text{reg}} \rangle} \geq \frac{\dim(M_{k-i(k)-n_0}(\Gamma))}{k}. \quad (1)$$

For ρ and k as in the Theorem, in [4] Shimura showed that

$$\lim_{k \rightarrow \infty} \frac{\dim(M_{k-i}(\Gamma))}{k} = c \quad (2)$$

and

$$\lim_{k \rightarrow \infty} \frac{\dim(M_k(\Gamma_1))}{k} = c\mu. \quad (3)$$

From (1) and (2) we easily deduce that

$$\liminf_{k \rightarrow \infty} \frac{\langle \rho, \rho_k \rangle}{k} \geq c \langle \rho, \rho_{\text{reg}} \rangle. \quad (4)$$

It is obvious that (3) is equivalent to

$$\lim_{k \rightarrow \infty} \sum_{\rho \in \mathcal{S}} \dim(\rho) \frac{\langle \rho, \rho_k \rangle}{k} = \sum_{\rho \in \mathcal{S}} \dim(\rho) c \langle \rho, \rho_{\text{reg}} \rangle.$$

Combining this equality with (4), we get

$$\lim_{k \rightarrow \infty} \frac{\langle \rho, \rho_k \rangle}{k} = c \langle \rho, \rho_{\text{reg}} \rangle,$$

as desired. $\square$

Acknowledgement. I would like to thank Professor David E. Rohrlich for his kindly help in revising this paper.

References

- [1] O. Forster, Lectures on Riemann Surfaces. Springer-Verlag, New York Inc., 1981.
- [2] N. Koblitz, Introduction to Elliptic Curves and Modular Forms. Springer-Verlag, New York Inc., 1984.
- [3] J.P. Serre, Linear Representations of Finite Groups. Springer-Verlag, New York Inc., 1977.
- [4] G. Shimura, Introduction to the Arithmetic Theory of Automorphic Functions. Princeton University Press, Princeton, New Jersey, 1971.